Noname manuscript No.
(will be inserted by the editor)

New Superpotential in Conservation Laws in General Relativity

Jiri Adamek · Jan Novotny

arXiv:1606.04352v2 [gr-qc] 26 Sep 2016

Received: date / Accepted: date

Abstract This work refers to the new formula for the superpotential U_i^{kl} in conservation laws in general relativity satisfying the integral and differential conservation laws within the Schwarzschild metric. The new su-perpotential is composed of two terms $U_i^{kl} = M_i^{kl} + A_i^{kl}$. The M_i^{kl} is based on Møller's concept [C. Møller, Ann. Phys. (NY) 4, 347 (1958)] and it is a function of the metric g_{ik} and its first derivative only. The second term A_i^{kl} is the antisymmetric tensor density of weight +1 and it consists of higher derivatives of the metric g_{ik} . Although the new superpotential consists of higher derivatives of the metric g_{ik} it might bring a new evaluation of the conservative quantities in general relativity.

Keywords superpotential · conservation laws · energy-momentum complex

1 Introduction

The recent observation of gravitational waves [1] might increase further investigation of energy-momentum complex τ_i^k within the conservation laws in general relativity [2,3], which is based on the definition of the superpotential U_i^{kl} where $\tau_i^k = U_i^{kl}{}_{,l}$. The variety of energy-momentum complex τ_i^k based on a different concept like Landau-Lifshits [4,5], Einstein-Freud [6], Møller [6, 7,8], Mickevic [8] can be used to study the conservative quantities [9,10,11] or gravitational waves [12]. Therefore, we can also find a different formula of the superpotential U_i^{kl} [13,14,15,16]. The gravitational energy-momentum

Institute of Plasma Physics of the CAS
Za Slovankou 3, Prague, Czech Republic
Tel.: +420-6605-2961
E-mail: adamek@ipp.cas.cz

Faculty of Science, Masaryk University
Kotlarska 2, Brno, Czech Republic

Table 1 Variety of the energy momentum complex and its constants in the formula for the general superpotential

	Einstein-Freud	Landau-Lifshitz	Møller	Mickevic
c_1	1	1	0	0
c_2	1	1	0	0
c_3	1	1	2	1
α	0.5	1	0.5	0.5

problem has been also discussed within the teleparallel gravity [17]. However, a different construction of the superpotential U_i^{kl} causes a problem in conservation laws in general relativity and it raises the question what is the "right" superpotential? The energy-momentum complex must yield the total energy of the whole system equal to Mc^2 and it must be invariant with respect to the Lorentz transformation and independent on the local coordinate system. This defines the integral and differential conservation laws in general relativity. It is shown that none of the above-mentioned candidates of energy momentum complex can satisfy all conservation laws. Therefore, we introduce the novel concept of the superpotential, which is composed of a two parts $U_i^{kl} = M_i^{kl} + A_i^{kl}$. The first term M_i^{kl}, is based on the Møller's concept and it is a function of the metric g_{ik} and its first derivative only. The second term A_i^{kl} is more complex and it consists of higher derivatives of the metric. It must be noted, that the formula for A_i^{kl} was found using a deductive and intuitive method. However, the novel concept of a superpotential satisfying the integral and differential conservations laws can contribute to the understanding of the gravitational energy in general relativity. A standard classification of the known superpotentials is based on the formula of the general superpotential [14]

$$U_i^{kl} = \frac{1}{2\kappa}\frac{c_1}{\alpha}\left(-g\right)^{\alpha}{}_{,m}\left(g^{lm}\delta_i^k - g^{km}\delta_i^l\right) + \frac{c_2}{2\kappa}\left(-g\right)^{\alpha}\left(g^{lm}{}_{,m}\,\delta_i^k - g^{km}{}_{,m}\,\delta_i^l\right) + \frac{c_3}{2\kappa}\left(-g\right)^{\alpha} g_{ib,m}\left(g^{bl}g^{mk} - g^{bk}g^{ml}\right) \tag{1}$$

with constant $\kappa = 8\pi G c^{-4}$ and the gravitational constant G. The general superpotential is composed of a linear combination of three terms, which are functions of the metric g_{ik} and its first derivative. The U_i^{kl} consists of four constants c_1, c_2, c_3 and α, which can be chosen in order to define a different energy-momentum complex τ_i^k. The following table represents the values of all constants mentioned above with respect to a different energy-momentum complex.

2 Application of the integral and differential conservation laws in Schwarzschild metric

The general superpotential U_i^{kl} (1) is assumed to be the most proper candidate for the evaluation of the energy-momentum complex τ_i^k or the general relativity conservation integral quantity four-momentum P_i

$$P_i = \frac{1}{c} \oint\limits_{\partial V} U_i^{0\lambda} \mathrm{d}\sigma_{0\lambda} \tag{2}$$

The four unknown constants can be determined by using differential and integral conditions. There exist two integral conditions in conservation laws. The first one demands the total energy of the insular system ($\partial V \to \infty$) to be $E = Mc^2$ or $P_0 = Mc$, which can be expressed as

$$P_i = \frac{1}{c} \oint\limits_{\partial V \to \infty} U_i^{0\lambda} \mathrm{d}\sigma_{0\lambda} = Mc\delta_i^0 \tag{3}$$
$$\lambda = 1,2,3 \quad i = 0,1,2,3$$

The second integral conservation law requires the four-momentum be invariant with respect to the Lorentz transformation. This leads to the condition

$$\frac{1}{c} \oint\limits_{\partial V \to \infty} U_i^{k\lambda} \mathrm{d}\sigma_{0\lambda} = 0 \tag{4}$$
$$\lambda, \kappa = 1,2,3 \quad i = 0,1,2,3$$

If we assume a metric g_{ik} of an external Schwarzschild solution [18] with stationary gravitation field in spherical coordinating system $x^i = (ct, r, \varphi, \vartheta)$

$$g_{00} = \left(1 - \frac{2m}{r}\right),\ g_{11} = -\left(1 - \frac{2m}{r}\right)^{-1},\ g_{22} = -r^2 sin^2\vartheta,\ g_{33} = -r^2 \tag{5}$$

with $m = G \cdot M/c^2$ then both integral conservation laws, (3) and (4), using the general superpotential (1) can be expressed by two linear equations

$$c_1 - \frac{c_2}{2} + \frac{c_3}{2} = 1 \tag{6}$$
$$2c_1 - c_2 - c_3 = 0 \tag{7}$$

with three unknown constants c_1, c_2, c_3. The remaining coefficient α in (1) is a free parameter and its value is chosen to be 0.5. Finally, the third equation will be found by using a differential condition, which states that the evaluation of the four-momentum P_i must be independent of the spatial coordinate. Only the third term of the superpotential (1) is independent of the spatial coordinate. Then the values of constants c_1, c_2 must be equal to zero in order to satisfy the differential conservation law. Finally, we obtain from (6) $c_3 = 2$ and from (7) $c_3 = 0$. In this case, the system of our equations is insolvable and therefore equation (1) does not represent the proper formula of the superpotential. This is the main problem of conservation laws in general relativity.

3 The novel concept of superpotential

A novel form of the superpotential has been found under the assumption that also higher derivatives of the metric g_{ik} should be taken into account. We propose here the new superpotential

$$U_i^{kl} = M_i^{kl} + A_i^{kl} \tag{8}$$

where

$$M_i^{kl} = \frac{c_3}{2\kappa}\sqrt{-g}\, g_{ib,m}\left(g^{bl}g^{mk} - g^{bk}g^{ml}\right) \tag{9}$$

$$A_i^{kl} = \frac{c_4}{2\kappa}\sqrt{-g}\, C S_{,m}\left(g^{lm}\delta_i^k - g^{km}\delta_i^l\right) \tag{10}$$

$$C = (S)^{\alpha}\left(S_{,a}S_{,b}g^{ab}\right)^{\beta}, \quad \alpha = \frac{3}{2},\ \beta = -1 \tag{11}$$

$$S = R^{abcd}R_{abcd} \tag{12}$$

The new concept of the superpotential is based on a linear combination of two terms. The first term M_i^{kl}(9) is based on Møller's concept and it is a function of the metric g_{ik} and its first derivative. It corresponds to the third term in the general formula of the superpotential (1). The second term A_i^{kl}(10) is an antisymmetric tensor density of weight +1. It is composed of scalar quantity C (11) and the Reimann tensor scalar invariant S (12) [18]. Therefore, the calculation of the four-momentum P_i (2) will be independent of the coordinate system, which satisfies the differential condition. This is the main idea of adding term A_i^{kl}. The final formula of the new superpotential (8) was found by a rather intuitive way and it consists of a higher derivative of the metric g_{ik}. This would require a Langrangian of the gravitational field $\sqrt{-g}L\left(g_{ik,lm}\,;g_{ik,l}\,;g_{ik}\right)$ [18] with higher derivatives. Despite the previous argument being non-trivial [19], we would then be able to satisfy both integral conservation laws (3) and (4) by searching for constant c_3 and c_4 in (9) and (10), respectively. The scalars C and S [18] in spherical coordinates in the Schwarzschild metric g_{ik} (5), assuming the asymptotic approximation $r \to \infty$ giving

$$S = 48\frac{m^2}{r^6};\ C = -\frac{1}{36\sqrt{48}}\frac{r^5}{m} \tag{13}$$

Then the first integral conservation law (3) can be expressed as a linear equation

$$\frac{c_3}{2} - \frac{c_4}{2\sqrt{12}} = 1 \tag{14}$$

The second integral conservation law (4) is more complex and can be expressed [8] as

$$\oint\limits_{\partial V\to\infty} M_i^{k\lambda}\mathrm{d}\sigma_{0\lambda} + \oint\limits_{\partial V\to\infty} A_i^{k\lambda}\mathrm{d}\sigma_{0\lambda} = -c_3\frac{1}{3}Mc^2\delta_i^k - c_4\frac{1}{3\sqrt{12}}Mc^2\delta_i^k = 0, \tag{15}$$

which yields the second linear equation

$$c_3 + \frac{c_4}{\sqrt{12}} = 0 \tag{16}$$

The linear equations (14) and (16) provide the constants

$$c_3 = 1 \quad c_4 = -\sqrt{12} \tag{17}$$

It is also interesting that the total energy of the insular system ($\partial V \to \infty$) as calculated in (14) is divided in two identical parts described by M_i^{kl} and A_i^{kl}. The final formula of the new superpotential is

$$U_i^{kl} = \frac{\sqrt{-g}}{2\kappa}\left[g_{ib,m}\left(g^{bl}g^{mk} - g^{bk}g^{ml}\right) + \sqrt{12}\,CS_{,m}\left(g^{km}\delta_i^l - g^{lm}\delta_i^k\right)\right] \tag{18}$$

4 Conclusions

It was shown that none of the linear combinations of the three terms in the general superpotential can yield a superpotential which satisfies all differential and integral conditions in the case of the Schwarzschild metric. In this paper we introduce a novel concept of the superpotential, which is based on the linear combination of Møller's concept M_i^{kl}, consisting only of the first derivative of the metric g_{ik}, and an additional term A_i^{kl} with a higher derivative of the metric g_{ik}. It seems that the presence of higher derivatives of the metric in the superpotential is necessary in order to satisfy the differential and both integral conditions. This would require a Langrangian of the gravitational field, which includes derivatives of the metric g_{ik} higher than second order.

Acknowledgements I would like thank to Dr. Jana Jurmanova and Dr. Jan Stöckel for the fruitful discussion and to Dr. Vladimir Fuchs for correcting English and to my family supporting me in my work.

References

1. B. P. Abbott et al., Phys. Rev. Lett. 116, 061102 (2016)
2. J. Horsky and J. Novotny, Czechoslovak Journal of Physics B, 19, 419-442 (1969)
3. S. Chandrasekhar, Astrophysical Journal, 158, page 45 (1969)
4. L. D. Landau, E. M. Lifshitz, The Classical Theory of Fields (1985)
5. K. S. Virbhadra , Physics Letters A, Volume 157, Issues 4-5, 195-197 (1991)
6. C. Møller, Ann. Phys. (NY) 4, 347 (1958)
7. J. Novotny, General relativity and gravitation 19, 1043 -1051 (1987)
8. C. Møller, Ann. Phys. 12, 118 (1961)
9. M. Favata, Phys. Rev. D 63, 064013 (2001)
10. K. S. Virbhadra and J. C. Parikh, Physics Letters B 317, 312-314 (1993)
11. G. G. L. Nashed, Il Nuovo Cimento B, 10, pp. 967-974 (2004)
12. D. Su and Y. Zhang, Phys. Rev. D 85, 104012 (2012)
13. O. Stolin and J. Novotny, Reports on Mathematical Physics, 48, 0034-4877 (2001)
14. J. Novotny, Folia Fac. Sci. Nat. UJEP Brunensis (Physica) 19, 1-55 (1974)
15. A. Komar, Phys. Rev. 113, 934 (1959)

16. M. Francavigli and M. Ferraris, Nuclear Physics B (Proc. Suppl.) 6, 405-407 (1989)
17. V. C. de Andrade, L. C. T. Guillen, and J. G. Pereira, Phys. Rev. Lett. 84, 4533-4536 (2000)
18. Ray d'Inverno, Introducing Einsteins relativity, Oxford University Press (1992)
19. S. V. Babak and L. P. Grishchuk, Phys. Rev. D 61, 024038 (1999)